

The most important thing to remember is that you must wash your hands thoroughly before handling any fresh piercing. The following recommendations are specific to Microdermals.

1. Clean your piercing once a day using a non-scented anti-bacterial soap such as Hibiclens®, Provon® or Dial®. Your piercing should only be cleaned once per day. For the first 3 weeks it must be pressure wrapped with a padded adhesive medical strip and from then on always cover with a band-aid while sleeping. Change the bandage when cleaning.
2. Clean your jewelry first by removing any secretions with a cotton swab and water. Use bottled water if you have well water.
3. Use soap on a cotton swab to remove loose and moist matter remaining on the piercing. Always use separate swabs for cleaning each side of the piercing. Never rotate the swab into the piercing – always rotate and wipe away from the piercing. Never use a cotton swab twice as you can swipe bacteria or other unwanted material into the piercing.
4. With a clean cotton swab remove any remaining cleaning agent. Wet the swab if necessary to remove any residue which may cause skin irritation or dryness. Tighten your beads if needed.
4. Pat dry with a clean, lint free paper towel. Never use cloth towels as they may hold bacteria. Never rub. Always keep it dry if not currently cleaning. Always remember to Pressure Wrap!!!!

Use cotton swabs to remove any lymph or secretions that may be present. Keep a 1 inch area surrounding your piercing completely clean of any make-up, lotion, scented and specialty soaps, or other similar product. Try to avoid smoky areas or smoking as much as possible. Avoid alcohol, aspirin, and any drugs not prescribed to you. A high zinc multivitamin, good hydration and a healthy sleep pattern will improve your healing overall. Make sure you are using clean sheets and pillow cases. Avoid pools, ponds, Jacuzzis, etc. Avoid others that may be sick, and wash your hands to avoid getting sick. Do not remove your jewelry unless instructed to by a professional piercer or your health care professional.

Consult a health care professional at the first sign of infection.

Symptoms of an infection can include: fever, redness, particularly if it spreads or forms a red streak, swelling or heat in the affected area, excessive pain or tenderness, and/or pus draining from or around the piercing. Follow all aftercare to reduce the risk of infections. Some people have sensitivities to metals and the appearance of a metal allergy and an infection are often very similar. If your piercing is in constant discomfort, has a discharge, is reddened beyond the first week without a causable factor, or simply doesn't feel "right" consult a health care professional. If a health care professional is unable to help or says that it is an allergic reaction and you need different jewelry, we are always glad to assist.

Your microdermal base is made of Grade 23 Titanium. The base will not need to be removed for MRI's. The tops are made from either 316LVM Stainless Steel or Grade 23 Titanium and should be removed for safety in any procedure. We can assist in removal of the top if necessary.

Purchase jewelry that is made of the same material. Do not put in any other jewelry until your piercing is healed. When it is healed and you wish to change the jewelry, sanitize the jewelry using isopropyl alcohol of 70% or greater strength. Allow to dry thoroughly. Clean the piercing as previously stated in steps 1 through 5 and then insert the new jewelry. If you wish to have assistance with jewelry removal or insertion, please call or stop in at Skinlab.

Do not touch your new piercing unless you are cleaning it!

Piercer _____

Date _____